

citrus



- Improves rooting of nursery trees
- Stimulates root and shoot development after establishment
- Better fruit set of bearing trees
- Better fruit mass
- Better yields
- Improves the development of new roots in spring
- Quicker recovery of trees after harvest




Kelpak, a natural biostimulant extracted from freshly harvested *Ecklonia maxima* kelp, scientifically proven to increase the health, quality and yield in a wide variety of crops. The global leader in auxin based seaweed products for over thirty years



Kelpak on Satsuma – Japan

3 x foliar sprays



**DROP FOR DROP
THE MOST EFFECTIVE
BIOSTIMULANT**

Summary of 5 soft skin citrus trials - South Africa 3 x foliar sprays				
TREATMENT	Fruits/tree (number)	Fruit mass (g/fruit)	Yield (kg/tree)	Increase (%)
CONTROL	390	130	45	-
KELPAK	400	140	52	+ 16%

RECOMMENDED APPLICATION RATE

Established trees: 0.3% Kelpak foliar spray applied at full bloom, and twice more with 2 to 3 week intervals for bigger fruit

Post harvest spray at 0.2% with nitrogen application and repeated 14 days later for recovery of heavy fruit bearing

Tree establishment: Apply 1 L of a 1.0% Kelpak solution (1 L Kelpak in 100 L water) as a soil drench directly after plant or drench the nursery bag with the solution directly before plant-out

For better tree development apply foliar sprays of 200 ml Kelpak per 100 L water (0.2%) with 3 to 4 week intervals during early growth

