


# strawberries



- Improved rooting
- Better shoot development after establishment
- Increased photosynthesis
- More marketable fruit
- Extended harvesting period
- Better yields



Kelpak, a natural biostimulant extracted from freshly harvested *Ecklonia maxima* kelp, scientifically proven to increase the health, quality and yield in a wide variety of crops. The global leader in auxin based seaweed products for over thirty years



Kelpak on strawberries globally				
COUNTRY	NUMBER OF TRIALS	RATE (L/ha)	FIRST APPLICATION	AVE. YIELD INCREASE
SOUTH AFRICA	6	3 L x 5*	At plant	22%
FRANCE	1	3 L x 5*	At plant	10%
HUNGARY	1	3 L x 3*	Start of spring growth	9%**
POLAND	2 seasons	3 L x 6*	At plant	23%

\*Applications normally 21 days apart


\*\* First 3 pickings



**DROP FOR DROP  
THE MOST EFFECTIVE  
BIOSTIMULANT**



Cumulative marketable yield per harvest week "Camerosa"



**RECOMMENDED APPLICATION RATE**

Dip runners/explants in 1% Kelpak solution before plant-out

Apply foliar spray at 3 L/ha, 3 weeks after plant-out

Repeat foliar spray 3 to 4 times, with 3 week intervals

**KELPAK**